arXiv:1102.1078v5 [math.CA] 31 Aug 2011

ON GENERALIZED COMPLETE ELLIPTIC INTEGRALS AND MODULAR FUNCTIONS

B. A. BHAYO AND M. VUORINEN

Abstract. This paper deals with generalized elliptic integrals and generalized modular functions. Several new inequalities are given for these and related functions.

2010 Mathematics Subject Classification: 33C99, 33B99
Keywords and phrases: Modular equation, generalized elliptic integral.

1. Introduction

During the past fifteen years, after the publication of the landmark paper [BBG], numerous papers have been written about generalized elliptic integrals, modular functions and inequalities for them. See e.g. [AQVV, AQ, B4, B5, HLVV, HVV, WZC, WZQC, ZWC1, ZWC2]. Modular equations have a long history, which goes back to the works of A.M. Legendre, K.F. Gauss, C. Jacobi and S. Ramanujan about number theory. Modular equations also occur in geometric function theory as shown in [AQVV, Vu2, K, LV] and in numerical computations of moduli of quadrilaterals [HRV]. For recent surveys of this topic from the point of view of geometric function theory, see [AVV4, AVV5, PV]. The study of these functions is motivated by potential applications to geometric function theory and to number theory.

Given complex numbers a, b and c with $c \neq 0, -1, -2, \ldots$, the *Gaussian hypergeometric function* is the analytic continuation to the slit place $\mathbf{C} \setminus [1, \infty)$ of the series

$$F(a,b;c;z) = {}_2F_1(a,b;c;z) = \sum_{n=0}^{\infty} \frac{(a,n)(b,n)}{(c,n)} \frac{z^n}{n!}, \qquad |z| < 1.$$

Here $(a,0) = 1$ for $a \neq 0$, and (a,n) is the *shifted factorial function* or the *Appell symbol*

$$(a,n) = a(a+1)(a+2)\cdots(a+n-1)$$

for $n \in \mathbb{Z}_+$.

`File: gcei110831.tex, printed: 2013-1-11, 21.40`

For later use we define classical *gamma function* $\Gamma(x)$, and *beta function* $B(x,y)$. For $\mathrm{Re}\, x>0$, $\mathrm{Re}\, y>0$, these functions are defined by

$$\Gamma(x)=\int_0^\infty e^{-t}t^{x-1}\,dt,\;\; B(x,y)=\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)},$$

respectively.

For the formulation of our main results and for later use we introduce some basic notation. The decreasing homeomorphism $\mu_a:(0,1)\to(0,\infty)$ is defined by

$$\mu_a(r)=\frac{\pi}{2\,\sin(\pi\, a)}\frac{F(a,1-a;1;r'^2)}{F(a,1-a;1;r^2)}=\frac{\pi}{2\,\sin(\pi\, a)}\frac{\mathcal{K}_a(r')}{\mathcal{K}_a(r)}$$

for $r\in(0,1)$ and $r'=\sqrt{1-r^2}$. A generalized modular equation with signature $1/a$ and order (or degree) p is

$$\mu_a(s)=p\,\mu_a(r),\quad 0<r<1. \tag{1.1}$$

We denote

$$s=\varphi^a_K(r)\equiv\mu_a^{-1}(\mu_a(r)/K),\quad K\in(0,\infty),\, p=1/K\,, \tag{1.2}$$

which is the solution of (1.1).

For $a\in(0,1/2]$, $K\in(0,\infty)$, $r\in(0,1)$, we have by [AQVV, Lemma 6.1]

$$\varphi^a_K(r)^2+\varphi^a_{1/K}(r')^2=1. \tag{1.3}$$

For $a\in(0,1/2]\,$, $r\in(0,1)$ and $r'=\sqrt{1-r^2}$, the generalized elliptic integrals are defined by

$$\left\{\begin{array}{l}\mathcal{K}_a(r)=\dfrac{\pi}{2}\,F(a,1-a;1;r^2),\\ \mathcal{K}'_a(r)=\,\mathcal{K}_a(r'),\\ \mathcal{K}_a(0)=\dfrac{\pi}{2},\;\mathcal{K}_a(1)=\infty,\end{array}\right.\quad\text{and}\quad\left\{\begin{array}{l}\mathcal{E}_a(r)=\dfrac{\pi}{2}\,F(a-1,1-a;1;r^2),\\ \mathcal{E}'_a(r)=\,\mathcal{E}_a(r'),\\ \mathcal{E}_a(0)=\dfrac{\pi}{2},\;\mathcal{E}_a(1)=\dfrac{\sin(\pi\,a)}{2(1-a)}.\end{array}\right.$$

In this paper we study the modular function $\varphi^a_K(r)$ for general $a\in(0,\frac12]$, as well as related functions $\mu_a,\ \mathcal{K}_a,\ \eta^a_K,\ \lambda_a$, and their dependency on r and K, where

$$\eta^a_K(x)=\left(\frac{s}{s'}\right)^2,\, s=\varphi^a_K(r),\, r=\sqrt{\frac{x}{1+x}}\,,\;\text{ for } x,K\in(0,\infty),$$

and

$$\lambda_a(K)=\left(\frac{\varphi^a_K(1/\sqrt2)}{\varphi^a_{1/K}(1/\sqrt2)}\right)^2=\left(\frac{\mu_a^{-1}(\pi/(2K\sin(\pi\,a))}{\mu_a^{-1}(\pi K/(2\sin(\pi\,a))}\right)^2=\eta^a_K(1). \tag{1.4}$$

Motivated by [L] and [BV] we define for $p>1$ and $r\in(0,1)$,

$$\mathrm{artanh}_p(x)=\int_0^x(1-t^p)^{-1}dt=xF\left(1\,,\frac1p;1+\frac1p;x^p\right)\,.$$

Then $\mathrm{artanh}_2(x)$ is the usual inverse hyperbolic tangent (artanh) function.

We give next some of the main results of this paper.

1.5. **Theorem.** *For* $a,b,c>0$, *and* $r\in(0,1)$, *the function* $g(p)=F(a,b;c;r^p)^{1/p}$ *is decreasing in* $p\in(0,\infty)$. *In particular, for* $p\geq 1$

$$F(a,b;c;r^p)^{1/p}\leq F(a,b;c;r)\leq F(a,b;c;r^{1/p})^p\,, \tag{1}$$

$$\left(\frac{\pi}{2}\right)^{1-1/p}\mathcal{K}_a(r^p)^{1/p}\leq \mathcal{K}_a(r)\leq\left(\frac{\pi}{2}\right)^{1-p}\mathcal{K}_a(r^{1/p})^p\,, \tag{2}$$

$$\left(\frac{\pi}{2}\right)^{1-p}\mathcal{E}_a(r^{1/p})^p\leq \mathcal{E}_a(r)\leq\left(\frac{\pi}{2}\right)^{1-1/p}\mathcal{E}_a(r^p)^{1/p}. \tag{3}$$

H. Alzer and S.-L. Qiu have given the following bounds for $\mathcal{K}=\mathcal{K}_{1/2}$ in [AQ, Theorem 18]

$$\frac{\pi}{2}\left(\frac{\operatorname{artanh}(r)}{r}\right)^{3/4}<\mathcal{K}(r)<\frac{\pi}{2}\left(\frac{\operatorname{artanh}(r)}{r}\right). \tag{1.6}$$

In the following theorem we generalize their result to the case of $\mathcal{K}_a$, and for the particular case $a=1/2$ our upper bound is better than their bound in (1.6). For a graphical comparison of the bounds see Figure 1 below.

1.7. **Theorem.** *For* $p\geq 2$ *and* $r\in(0,1)$, *we have*

$$\begin{aligned}\frac{\pi}{2}\left(\frac{\operatorname{artanh}_p(r)}{r}\right)^{1/2} &< \frac{\pi}{2}\left(1-\frac{p-1}{p^2}\log(1-r^2)\right)\\ &< \mathcal{K}_a(r)<\frac{\pi}{2}\left(1-\frac{2}{p\,\pi_p}\log(1-r^2)\right),\end{aligned}$$

where $a=1/p$ *and* $\pi_p=2\pi/(p\sin(\pi/p))$.

In [AQVV, Theorem 5.6] (see also [BPV, Theorem 1.5, 1.8]) it was proved that for $a\in(0,1/2]$ we have

$$\mu_a\left(\frac{r\,s}{1+r's'}\right)\leq\mu_a(r)+\mu_a(s)\leq 2\mu_a\left(\frac{\sqrt{2rs}}{\sqrt{1+r\,s+r'\,s'}}\right),$$

for all $r,s\in(0,1)$. This inequality will be generalized below in Theorem 4.3. In the next theorem we give a similar result for the function $\mathcal{K}_a$.

1.8. **Theorem.** *The function* $f(x)=1/\mathcal{K}_a(1/\cosh(x))$ *is increasing and concave from* $(0,\infty)$ *onto* $(0,2/\pi)$. *In particular,*

$$\frac{\mathcal{K}_a(r)\,\mathcal{K}_a(s)}{\mathcal{K}_a(rs/(1+r's'))}\leq\mathcal{K}_a(r)+\mathcal{K}_a(s)\leq\frac{2\,\mathcal{K}_a(r)\,\mathcal{K}_a(s)}{\mathcal{K}_a(\sqrt{rs/(1+rs+r's')})}\leq\frac{2\,\mathcal{K}_a(r)\,\mathcal{K}_a(s)}{\mathcal{K}_a(rs)},$$

for all $r,s\in(0,1)$, *with equality in the third inequality if and only if* $r=s$.

There are several bounds for the function $\mu_a(r)$ when $a=1/2$ in [AVV1, Chap.5]. In the next theorem we give a twosided bound for $\mu_a(r)$.

1.9. **Theorem.** *For* $p \geq 2$ *and* $r \in (0,1)$*, let*

$$l_p(r) = \left(\frac{\pi_p}{2}\right)^2 \left(\frac{p^2-(p-1)\log r^2}{p\,\pi_p - 2\log r'^2}\right) \quad \text{and} \quad u_p(r) = \left(\frac{p}{2}\right)^2 \left(\frac{p\,\pi_p - 2\log r^2}{p^2-(p-1)\log r'^2}\right).$$

(1) The following inequalities hold

$$l_p(r) < \mu_a(r) < u_p(r)\,,$$

where $a = 1/p$*.*
(2) For $p = 2$ *we have*

$$u_2(r) < \frac{4}{\pi}\, l_2(r)\,.$$

Acknowledgments. The first author is indebted to the Graduate School of Mathematical Analysis and its Applications for support. The second author was, in part, supported by the Academy of Finland, Project 2600066611. Both authors wish to acknowledge the expert help of Dr. H. Ruskeepää in the use of the Mathematica® software [Ru]. The authors also acknowledge the constructive suggestions of the referee.

2. Proofs of Theorems 1.5,1.7,1.8 and 1.9.

For easy reference we record the next two lemmas from [AVV1] which have found many applications. Some of the applications are reviewed in [AVV5]. The first result sometimes called the *monotone l'Hospital rule.*

2.1. **Lemma.** [AVV1, Theorem 1.25] *For* $-\infty < a < b < \infty$*, let* $f, g : [a,b] \to \mathbb{R}$ *be continuous on* $[a,b]$*, and be differentiable on* (a,b)*. Let* $g'(x) \neq 0$ *on* (a,b)*. If* $f'(x)/g'(x)$ *is increasing (decreasing) on* (a,b)*, then so are*

$$[f(x)-f(a)]/[g(x)-g(a)] \quad \text{and} \quad [f(x)-f(b)]/[g(x)-g(b)].$$

If $f'(x)/g'(x)$ *is strictly monotone, then the monotonicity in the conclusion is also strict.*

2.2. **Lemma.** [AVV1, Lemma 1.24] *For* $p \in (0,\infty]$*, let* $I = [0,p)$*, and suppose that* $f, g : I \to [0,\infty)$ *are functions such that* $f(x)/g(x)$ *is decreasing on* $I \setminus \{0\}$ *and* $g(0) = 0$ *and* $g(x) > 0$ *for* $x > 0$*. Then*

$$f(x+y)(g(x)+g(y)) \leq g(x+y)(f(x)+f(y))\,,$$

for $x, y, x+y \in I$*. Moreover, if the monotonicity of* $f(x)/g(x)$ *is strict then the above inequality is also strict on* $I \setminus \{0\}$*.*

For easy reference we recall the following lemmas from [AQVV].

2.3. **Lemma.** *For* $a \in (0,1/2]$*,* $K \in (1,\infty)$*,* $r \in (0,1)$ *and* $s = \varphi_K^a(r)$*, we have*

(1) $f(r) = s'\,\mathcal{K}_a(s)^2/(r'\,\mathcal{K}_a(r)^2)$ *is decreasing from* $(0,1)$ *onto* $(0,1)$*,*
(2) $g(r) = s\,\mathcal{K}_a'(s)^2/(r\,\mathcal{K}_a'(r)^2)$ *is decreasing from* $(0,1)$ *onto* $(1,\infty)$*,*

(3) *the function* $r'^c\,\mathcal{K}_a(r)$ *is decreasing if and only if* $c \geq 2a(1-a)$*, in which case* $r'^c\,\mathcal{K}_a(r)$ *is decreasing from* $(0,1)$ *onto* $(0,\pi/2)$*. Moreover,* $\sqrt{r'}\,\mathcal{K}_a(r)$ *is decreasing for all* $a \in (0,1/2]$*.*

2.4. **Lemma.** *The following formulae hold for* $a \in (0,1/2]$, $r \in (0,1)$ *and* $x, y, K \in (0,\infty)$*:*

(1) $\dfrac{dF}{dr} = \dfrac{l\,m}{n} F(1+l, 1+m; 1+n; r);\ F = F(l,m;n;r)\,,$

(2) $\dfrac{d\,\mathcal{K}_a(r)}{dr} = \dfrac{2(1-a)(\,\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r))}{rr'^2}\,,$

(3) $\dfrac{d\,\mathcal{E}_a(r)}{dr} = \dfrac{2(a-1)(\,\mathcal{K}_a(r) - \,\mathcal{E}_a(r))}{r}\,,$

(4) $\dfrac{d\mu_a(r)}{dr} = \dfrac{-\pi^2}{4rr'^2\,\mathcal{K}_a(r)^2}\,,$

(5) $\dfrac{d\varphi_K^a(r)}{dr} = \dfrac{ss'^2\,\mathcal{K}_a(s)^2}{Krr'^2\,\mathcal{K}_a(r)^2} = \dfrac{ss'^2\,\mathcal{K}_a(s)\,\mathcal{K}'_a(s)}{rr'^2\,\mathcal{K}_a(r)\,\mathcal{K}'_a(r)} = K\dfrac{ss'^2\,\mathcal{K}'_a(s)^2}{rr'^2\,\mathcal{K}'_a(r)^2}\,,$

(6) $\dfrac{d\varphi_K^a(r)}{dK} = \dfrac{4ss'^2\,\mathcal{K}_a(s)^2\mu_a(r)}{\pi^2K^2}\,,$
where $s = \varphi_K^a(r)$*,*

(7) $\dfrac{d\eta_K^a(x)}{dx} = \dfrac{1}{K}\left(\dfrac{r'\,s\,\mathcal{K}_a(s)}{rs'\,\mathcal{K}_a(r)}\right)^2 = K\left(\dfrac{r'\,s\,\mathcal{K}'_a(s)}{rs'\,\mathcal{K}'_a(r)}\right)^2 = \left(\dfrac{r'\,s}{rs'}\right)^2 \dfrac{\mathcal{K}_a(s)\,\mathcal{K}'_a(s)}{\mathcal{K}_a(r)\,\mathcal{K}'_a(r)}\,,$

(8) $\dfrac{d\eta_K^a(x)}{dK} = \dfrac{8\eta_K^a(x)\mu_a(r)K_a(s)^2}{\pi^2K^2}\,,$
in (7) *and* (8)*,* $r = \sqrt{x/(1+x)}$ *and* $s = \varphi_K^a(r)$*.*

2.5. **Lemma.** [AVV1, Theorem 1.52(1)] *For* $a, b > 0$*, the function*

$$f(x) = \frac{F(a,b;a+b;x) - 1}{\log(1/(1-x))}$$

is strictly increasing from $(0,1)$ *onto* $(ab/(a+b), 1/B(a,b))$*.*

2.6**. Proof of Theorem 1.5.** With $G(r) = F(a,b;c;r^p)$, and g as in Theorem 1.5 we get by Lemma 2.4(1)

$$g'(p) = -\frac{(G(r))^{1/p-1}}{cp^2}\left(c\,G(r)\log\left(\,G(r)\right) + a\,b\,p\,r^p\,F\left(a+1, b+1; c+1; r^p\right)\log(1/r)\right)$$

which is negative. Hence this implies (1), and (2) follows from (1). For (3), write $F(r) = F(-a,b;c;r^p)$. We define $h(p) = F(r)^{1/p}$ and get

$$h'(p) = \frac{(F(r))^{1/p-1}}{cp^2}\left(c\,F(r)\log\left(1/\,F(r)\right) + a\,b\,p\,r^p\,F\left(a+1, b+1; c+1; r^p\right)\log(1/r)\right)$$

which is positive because $F(r) \in (0,1)$. Hence h is increasing in p, and (3) follows easily. $\square$



FIGURE 1. Comparison of upper bounds given in Theorem 1.7 and (1.6) for $\mathcal{K}(r)$.

2.7. **Proof of Theorem 1.7.** By the definition of $\mathrm{artanh_p}$, Lemma 2.5 and Bernoulli inequality we get

$$\begin{aligned}\left(\frac{\mathrm{artanh}_p(r)}{r}\right)^{1/2} &= \left(F\left(1,\frac{1}{p};1+\frac{1}{p};r^p\right)\right)^{1/2}\\ &< \left(1-\frac{1}{p}\log(1-r^p)\right)^{1/2}\\ &\le 1+\frac{1}{2p}\log\left(\frac{1}{1-r^p}\right)\\ &\le 1+\frac{p-1}{p^2}\log\left(\frac{1}{1-r^p}\right)\\ &\le 1-\frac{p-1}{p^2}\log(1-r^2)=\xi.\end{aligned}$$

Again by Lemma 2.5 and [AS, 6.1.17] we obtain

$$\begin{aligned}\xi &< F\left(\frac{1}{p},1-\frac{1}{p};1;r^2\right)=\frac{2}{\pi}\mathcal{K}_{1/p}(r)\\ &< 1-\frac{1}{B(1/p,1-1/p)}\log(1-r^2)\\ &= 1-\frac{2}{p\,\pi_p}\log(1-r^2)\,,\end{aligned}$$

and this completes the proof. $\square$

2.8. **Proof of Theorem 1.8.** Writing $r = 1/\cosh(x)$ we have

$$\frac{dr}{dx} = -(\sinh x)/\cosh^2 x = -r\,r'$$

and

$$\begin{aligned} f'(x) &= -\frac{\mathcal{K}_a'(r)}{\mathcal{K}_a^2(r)}\frac{dr}{dx} = -\frac{2(1-a)}{\mathcal{K}_a^2(r)}\,\frac{\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r)}{r\,r'^2}\,(-r\,r') \\ &= 2(1-a)\frac{\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r)}{r'\;\mathcal{K}_a(r)^2}\,, \end{aligned}$$

which is positive and increasing in r by Lemma 2.3(3) and therefore $f'(x)$ is decreasing in x and f is concave. Hence,

$$\begin{aligned} &\frac{1}{2}(f(x)+f(y)) \le f\left(\frac{x+y}{2}\right) \\ \iff &\frac{1}{2}\left(\frac{1}{\mathcal{K}_a(1/\cosh(x))} + \frac{1}{\mathcal{K}_a(1/\cosh(y))}\right) \le \frac{1}{\mathcal{K}_a(1/\cosh((x+y)/2))} \\ \iff &\mathcal{K}_a(r) + \mathcal{K}_a(s) \le \frac{2\,\mathcal{K}_a(r)\,\mathcal{K}_a(s)}{\mathcal{K}(\sqrt{rs/(1+rs+r^{'}s^{'})})}\,, \end{aligned}$$

using $\cosh^2((x+y)/2) = (1+rs+r^{'}s^{'})/(rs)$ and setting $s = 1/\cosh(y)$. Clearly,

$$(r-s)^2 \ge 0 \iff 1 - 2rs + r^2s^2 \ge 1 - r^2 - s^2 + r^2s^2$$

$$\iff 1 - rs \ge r^{'}s^{'} \iff 2 \ge 1 + rs + r^{'}s^{'} \iff 2rs/(1+rs+r^{'}s^{'}) \ge rs\,,$$

and the third inequality follows. Obviously, $f(0+) = 0$, and $f^{'}(x)$ is decreasing in x. Then $f(x)/x$ is decreasing and $f(x+y) \le f(x) + f(y)$ by Lemmas 2.1 and 2.2, respectively. This implies the first inequality.

2.9. **Proof of Theorem 1.9.** By Lemma 2.5 we get

(a) $1 - \dfrac{p-1}{p^2}\log r^2 < F\left(\dfrac{1}{p}, 1-\dfrac{1}{p}; 1; 1-r^2\right) < 1 - \dfrac{2}{p\,\pi_p}\log r^2$

(b) $1 - \dfrac{p-1}{p^2}\log(1-r^2) < F\left(\dfrac{1}{p}, 1-\dfrac{1}{p}; 1; r^2\right) < 1 - \dfrac{2}{p\,\pi_p}\log(1-r^2)\,.$

By using (a), (b) and the definition of μ_a, we get (1). The claim (2) is equivalent to

$$\frac{2(\pi - \log(r^2))}{4 - \log(1-r^2)} < \frac{4}{\pi}\left(\frac{\pi}{2}\right)^2 \frac{4 - \log(r^2)}{\pi - \log(1-r^2)}$$

$$\iff 4(\pi - \log(r^2))(\pi - \log(1-r^2)) - (4 - \log(r^2))(4 - \log(1-r^2)) < 0,$$

$$\iff (\pi - 4)(4\pi - \log(r^2)\log(1-r^2)) < (\pi - 4)(4\pi - (\log(2))^2) < 0\,.$$

For the second last inequality we define $w(x) = \log(x)\log(1-x)$, and get

$$w^{'}(x) = \frac{(1-x)\log(1-x) - x\log(x)}{x(1-x)} = \frac{-g(x)}{x(1-x)}\,,$$

and see that $g(x) = x\log(x) - (1-x)\log(1-x)$ is convex on $(0,1/2)$ and concave on $(1/2,1)$. This implies that $g(x) < 0$ for $x \in (0,1/2)$ and $g(x) > 0$ for $x \in (1/2,1)$. Therefore w is increasing in $(0,1/2)$ and decreasing in $(1/2,1)$. Hence the function w has a global maximum at $x = 1/2$ and this completes the proof.

□

One can obtain the following inequalities by using the proof of Theorem 1.9:

$$\frac{p\,\pi_p}{2\pi}\frac{\mathcal{K}_a(r)}{(1-(2/(p\,\pi_p))\log r^2)} \le \mu_a(r') \le \frac{p\,\pi_p}{2\pi}\frac{\mathcal{K}_a(r)}{(1-((p-1)/p)\log r^2)}\,,$$

with $a = 1/p$ and $p \ge 2$.

2.10. **Lemma.** *The following inequalities hold for all* $r, s \in (0,1)$ *and* $a \in (0,1/2]$*;*

(1) $\mathcal{K}_a(r\,s) \le \sqrt{\mathcal{K}_a(r^2)\,\mathcal{K}_a(s^2)} \le \frac{2}{\pi}\,\mathcal{K}_a(r)\,\mathcal{K}_a(s)\,,$

(2) $\frac{2}{\pi}\,\mathcal{E}_a(r)\,\mathcal{E}_a(s) \le \sqrt{\mathcal{E}_a(r^2)\,\mathcal{E}_a(s^2)} \le \mathcal{E}_a(rs)\,.$

Proof. Define $f(x) = \log(\mathcal{K}_a(e^{-x}))$, $x > 0$. We get by Lemma 2.4(2)

$$f'(x) = -2(1-a)\frac{\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r)}{r'^2\,\mathcal{K}_a(r)}\,, \quad r = e^{-x}\,,$$

and this is negative by the fact that $h(r) = \mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r) > 0$ and decreasing in r by [AQVV, Lemma 5.4(1)] and the fact that h is increasing ($h'(r) = 2ar\,\mathcal{K}_a(r) > 0$). Therefore $f'(x)$ is increasing in x, hence f is convex, and this implies the first inequality of part one. The second inequality follows from Theorem 1.5(2).

The first inequality of part two follows from the Theorem 1.5(3), for the second inequality we define $g(x) = \log(\mathcal{E}_a(z))$, $z = e^{-x}$, $x > 0$, and get Lemma 2.4(3)

$$g'(x) = 2(1-a)\frac{(\mathcal{K}_a(z) - \mathcal{E}_a(z))}{\mathcal{E}_a(z)}\,,$$

which is positive and increasing in z by [AQVV, Theorem 4.1(3), Lemma 5.2(3)], hence $g'(x)$ is decreasing in x, therefore g is increasing and concave. This implies that

$$\log(\mathcal{E}_a(e^{-(x+y)/2})) \ge (\log(\mathcal{E}_a(e^{-x})) + \log(\mathcal{E}_a(e^{-y})))/2\,,$$

and the second inequality follows if we set $r = e^{-x/2}$ and $s = e^{-y/2}$. □

3. FEW REMARKS ON SPECIAL FUNCTIONS

In this section we generalize some results from [AVV1, Chapter 10].

3.1. **Theorem.** *The function* $\mu_a^{-1}(y)$ *has exactly one inflection point and it is log-concave from* $(0,\infty)$ *onto* $(0,1)$*. In particular,*

$$(\mu_a^{-1}(x))^p(\mu_a^{-1}(y))^q \le \mu_a^{-1}(p\,x + q\,y)$$

for p, q, x, $y > 0$ *with* $p + q = 1$.

Proof. Letting $s = \mu_a^{-1}(y)$ we see that $\mu_a(s) = y$. By Lemma 2.4(4) we get

$$\frac{ds}{dy} = -\frac{4}{\pi^2} s s'^2 \mathcal{K}_a(s)^2 \,,$$

$$\begin{aligned}\frac{d^2 s}{dy^2} &= -\frac{ds}{dy}\frac{4}{\pi^2}(s'^2 \mathcal{K}_a(s)^2 - 2s^2 \mathcal{K}_a(s)^2 + 2\,\mathcal{K}_a(s)^2(\,\mathcal{E}_a(s) - s'^2 \mathcal{K}_a(s)))\\ &= \frac{16}{\pi^4} s s'^2 \mathcal{K}_a(s)^3 (2\,\mathcal{E}_a(s) - (1+s^2)\,\mathcal{K}_a(s))\,.\end{aligned}$$

We see that $2\,\mathcal{E}_a(s) - (1+s^2)\,\mathcal{K}_a(s)$ is increasing from $(0,\infty)$ onto $(-\infty, \pi/2)$ as a function of y. Hence $d(\mu_a^{-1}(y_0))/dy^2 = 0$, for $y_0 \in (0,\infty)$ and μ_a^{-1} has exactly one inflection point. Let $f(y) = \log(\mu_a^{-1}(y)) = \log\, s$. Then

$$f'(y) = -\frac{4}{\pi^2} s'^2 \mathcal{K}_a(s)^2$$

which is decreasing as a function of y, by Lemma 2.3(3), hence μ_a^{-1} is log-concave. This completes the proof. □

3.2. **Corollary.** (1) *For* $K \geq 1$, *the function* $f(r) = (\log \varphi_K^a(r))/\log r$ *is strictly decreasing from* $(0,1)$ *onto* $(0, 1/K)$.
(2) *For* $K \geq 1\,, r \in (0,1)$, *the function* $g(p) = \varphi_K^a(r^p)^{1/p}$ *is decreasing from* $(0,\infty)$ *onto* $(r^{1/K}, 1)$. *In particular,*

$$r^{p/K} \leq \varphi_K^a(r^p) \leq \varphi_K^a(r)^p,\ \ p \geq 1\,,$$

and

$$\varphi_K^a(r^p) \geq \varphi_K^a(r)^p, \quad 0 < p \leq 1\,.$$

Proof. Let $s = \varphi_K^a(r)$. By Lemma 2.4(5) we get

$$f'(r) = \frac{r s s'^2}{s r r'^2}\frac{\mathcal{K}_a(s)\,\mathcal{K}_a'(s)}{\mathcal{K}_a(r)\,\mathcal{K}_a'(r)} \log r - \log s,$$

and this is equivalent to

$$r(\log r)^2 f'(r) = s'^2\,\mathcal{K}_a(s)\,\mathcal{K}_a'(s)\left(\frac{\log r}{r'^2\,\mathcal{K}_a(r)\,\mathcal{K}_a'(r)} - \frac{\log s}{s'^2\,\mathcal{K}_a(s)\,\mathcal{K}_a'(s)}\right),$$

which is negative by Lemma 2.3(3). The limiting values follow from l'Hôpital Rule and Lemma 2.3(1). We observe that

$$\log g(p) = \left(\frac{\log \varphi_K^a(r^p)}{\log(r^p)}\right)\log r\,,$$

and (2) follows from (1). □

3.3. **Lemma.** *For* $0 < a \leq 1/2$, $K, p \geq 1$ *and* $r, s \in (0,1)$, *the following inequalities hold*

$$\frac{\sqrt[p]{\varphi_K^a(r^p)} + \sqrt[p]{\varphi_K^a(s^p)}}{1 + \sqrt[p]{\varphi_K^a(r^p)\varphi_K^a(s^p)}} \leq \frac{\varphi_K^a(r) + \varphi_K^a(s)}{1 + \varphi_K^a(r)\varphi_K^a(s)} \leq \frac{\varphi_K^a(\sqrt[p]{r})^p + \varphi_K^a(\sqrt[p]{s})^p}{1 + (\varphi_K^a(\sqrt[p]{r})\varphi_K^a(\sqrt[p]{s}))^p}\,.$$



FIGURE 2. Let $g(a,K,p,r,s) = \frac{\sqrt[p]{\varphi_K^a(r^p)}+\sqrt[p]{\varphi_K^a(s^p)}}{1+\sqrt[p]{\varphi_K^a(r^p)\varphi_K^a(s^p)}}$, $h(a,K,r,s) = \varphi_K^a\left(\frac{r+s}{1+rs}\right)$ be the lower bounds in Lemma 3.3 and (3.4), respectively. For $a = 0.2$, $K = 1.5$, $p = 1.3$ and $s = 0.5$ the functions g and h are plotted. We see that for $r \in (0.2, 1)$ the first lower bound is better.

Proof. It follows from the Corollary 3.2(2) that

$$\varphi_K^a(r^p)^{1/p} \le \varphi_K^a(r)\,.$$

From the fact that artanh is increasing, we conclude that

$$\operatorname{artanh}(\varphi_K^a(r^p)^{1/p}) + \operatorname{artanh}(\varphi_K^a(s^p)^{1/p}) \le \operatorname{artanh}(\varphi_K^a(r)) + \operatorname{artanh}(\varphi_K^a(s))\,.$$

This is equivalent to

$$\operatorname{artanh}\left(\frac{\varphi_K^a(r^p)^{1/p}+\varphi_K^a(s^p)^{1/p}}{1+(\varphi_K^a(r^p)+\varphi_K^a(s^p))^{1/p}}\right) \le \operatorname{artanh}\left(\frac{\varphi_K^a(r)+\varphi_K^a(s)}{1+(\varphi_K^a(r)+\varphi_K^a(s))}\right),$$

and the first inequality holds. Similarly, the second inequality follows from $\varphi_K^a(r) \le \varphi_K^a(r^{1/p})^p$. $\square$

For $0 < a \le 1$, $K \ge 1$ and $r, s \in (0,1)$, the following inequality

$$\varphi_K^a\left(\frac{r+s}{1+rs}\right) \le \frac{\varphi_K^a(r)+\varphi_K^a(s)}{1+\varphi_K^a(r)\varphi_K^a(s)} \tag{3.4}$$

is given in [AQVV, Remark 6.17].

3.5. **Theorem.** *For $r, s \in (0,1)$, we have*

(1) $|\varphi_K^a(r) - \varphi_K^a(s)| \le \varphi_K^a(|r-s|) \le e^{(1-1/K)R(a)/2}|r-s|^{1/K}$, $K \ge 1$, *here $R(a)$ is as in* [AQVV, Theorem 6.7]

(2) $|\varphi_K^a(r)-\varphi_K^a(s)| \geq \varphi_K^a(|r-s|) \geq e^{(1-1/K)R(a)/2}|r-s|^{1/K}$, $0<K\leq 1$.

Proof. It follows from [AQVV, Theorem 6.7] that $r^{-1}\varphi_K^a(r)$ is decreasing on $(0,1)$, if $K>1$ and by Lemma 2.2 we obtain

$$\varphi_K^a(x+y) \leq \varphi_K^a(x)+\varphi_K^a(y),\;\; x,y\in(0,1)\,.$$

Now the first inequality in (1) follows if we take $r=x+y$ and $s=y$, the second one follows from [AQVV, Theorem 6.7]. Next, (2) follows from (1) and the fact that

$$\varphi_{AB}^a(r)=\varphi_A^a(\varphi_B^a(r)),\;\; A,B>0\,, r\in(0,1)$$

when we replace K, r and s by $1/K$, $\varphi_{1/K}^a(r)$, $\varphi_{1/K}^a(s)$, respectively. ☐

3.6. **Theorem.** *For $a\in(0,1/2]$, $c,r\in(0,1)$ and $K,L\in(0,\infty)$ we have*

1. *The function $f(K)=\log(\varphi_K^a(r))$ is increasing and concave from $(0,\infty)$ onto $(-\infty,0)$.*
2. *The function $g(K)=\operatorname{artanh}(\varphi_K^a(r))$ is increasing and convex from $(0,\infty)$ onto $(0,\infty)$.*
3. $\varphi_K^a(r)^c\varphi_L^a(r)^{1-c} \leq \varphi_{cK+(1-c)L}^a(r) \leq \tanh(c\operatorname{artanh}(\varphi_K^a(r))+(1-c)\operatorname{artanh}(\varphi_L^a(r)))\,.$
4. $$\sqrt{\varphi_K^a(r)\varphi_L^a(r)} \leq \varphi_{(K+L)/2}^a(r)$$ $$\leq \frac{\varphi_K^a(r)+\varphi_L^a(r)}{1+\varphi_K^a(r)\varphi_L^a(r)+\varphi_{1/K}^a(r')\varphi_{1/L}^a(r')}\,.$$

Proof. For (1), by Lemma 2.4(6) we get

$$f'(K)=4s'^2\,\mathcal{K}_a(s)^2\mu_a(r)/(\pi^2K),$$

which is positive and decreasing by Lemma 2.3(3). For (2), we get

$$f'(K)=4s\,\mathcal{K}_a(s)^2\mu_a(r)/(\pi^2K^2)=s\,\mathcal{K}_a'(s)^2/\mu_a(r)$$

by Lemma 2.4(6), which is positive and increasing by Lemma 2.3(3). By (1) and (2) we get

$$c\log(\varphi_K^a(r))+(1-c)\log(\varphi_L^a(r)) \leq \log(\varphi_{cK+(1-a)L}^a(r)),$$

$$\operatorname{artanh}(\varphi_{cK+(1-c)K}^a(r)) \leq a\operatorname{artanh}(\varphi_K^a(r))+(1-c)\operatorname{artanh}(\varphi_L^a(r)),$$

respectively, and (3) follows. Also

$$(\log(\varphi_K^a(r))+\log(\varphi_L^a(r)))/2 \leq \log(\varphi_{(K+L)/2}^a(r))\,,$$

and

$$\operatorname{artanh}(\varphi_{(K+L)/2}^a(r)) \leq (\operatorname{artanh}(\varphi_K^a(r))+\operatorname{artanh}(\varphi_L^a))/2\,,$$

follow from (1) and (2), and hence (4) holds. ☐

3.7. **Theorem.** *For $K\geq 1$ and $0<m<n$, the following inequalities hold*

(1) $\eta_K^a(m\,n) \leq \sqrt{\eta_K^a(m^2)\eta_K^a(n^2)}\,,$

(2) $\left(\dfrac{n}{m}\right)^{1/K} < \dfrac{\eta_K^a(n)}{\eta_K^a(m)} < \left(\dfrac{n}{m}\right)^{K}$,

(3) $\eta_K^a(m)\eta_K^a(n) < \left(\eta_K^a\left(\dfrac{m+n}{2}\right)\right)^2$,

(4) $2\dfrac{\eta_K^a(m)\eta_K^a(n)}{\eta_K^a(m)+\eta_K^a(n)} < \eta_K^a(\sqrt{m\,n}) < \sqrt{\eta_K^a(m)\eta_K^a(n)}$.

Proof. We define a function $g(x) = \log \eta_K^a(e^x)$ on $\mathbb{R}$. By [AQVV, Theorem 1.16], g is increasing, convex and satisfies $1/K \le g'(x) \le K$. Then

$$\begin{aligned}\log \eta_K^a(e^{(x+y)/2}) &= g\left(\frac{x+y}{2}\right) \le \frac{g(x)+g(y)}{2}\\ &= \frac{1}{2}\log(\eta_K^a(e^x)) + \frac{1}{2}\log(\eta_K^a(e^y)),\end{aligned}$$

and this is equivalent to

$$\log \eta_K^a(e^{x/2}e^{y/2}) \le \log(\eta_K^a(e^{x/2})\eta_K^a(e^{y/2}))\,.$$

Hence (1) follows if we set $e^{x/2} = m$ and $e^{y/2} = n$. For (2), let $x > y$. Then by the inequality $1/K \le g'(x) \le K$ and the mean value theorem we get

$$(x-y)/K \le g(x) - g(y) \le K(x-y),$$

and this is equivalent to

$$(\log(e^x) - \log(e^y))/K \le \log(\eta_K^a(e^x)) - \log(\eta_K^a(e^y)) \le K(\log(e^x) - \log(e^y))\,.$$

By setting $e^{x/2} = m$ and $e^{y/2} = n$ we get the desired inequality. For (3), let $f(x) = \log(\eta_K^a(x))$, $r = \sqrt{x/(1+x)}$ and $s = \varphi_K^a(r)$. Then by Lemma 2.4(7) we get

$$\begin{aligned}f'(x) &= \frac{1}{K}\left(\frac{s'}{s}\right)^2\left(\frac{sr'\,\mathcal{K}_a(s)}{rs'\,\mathcal{K}_a(r)}\right)^2 = \frac{1}{K}\left(\frac{r'}{r}\right)^2\left(\frac{\mathcal{K}_a(s)}{\mathcal{K}_a(r)}\right)^2\\ &= \frac{1}{K}\left(\frac{r'}{s}\right)^2\left(\frac{s\,\mathcal{K}_a(s)}{r\,\mathcal{K}_a(r)}\right)^2,\end{aligned}$$

which is positive and decreasing by Lemma 2.3(2). Hence $(f(x)+f(y))/2 \le f((x+y)/2)$, and the inequality follows.

For (4), letting $h(x) = 1/\eta_K^a(e^x)$, we see that this is log-concave by (1), and we get

$$\frac{\log(1/\eta_K^a(e^x)) + \log(1/\eta_K^a(e^y))}{2} < \log(1/\eta_K^a(e^{(x+y)/2}))\,,$$

Setting $e^x = m$ and $e^y = n$ we get the second inequality. We observe that $h(x) = (s'/s)$, $s = \varphi_K^a(r)$, $r = \sqrt{e^x/(e^x+1)}$. We get

$$-f'(x) = \frac{1}{K}\left(\frac{r'}{s}\right)\left(\frac{s'\,\mathcal{K}_a(s)}{r'\,\mathcal{K}_a(r)}\right)^2,$$

which is positive and decreasing by Lemma 2.3(1), hence h is convex, and the first inequality follows easily. $\square$

3.8. **Theorem.** *For* $x \in (0,\infty)$, *the function* $f : (0,\infty) \to (0,\infty)$ *defined by* $f(K) = \eta_K^a(x)$ *is increasing, convex and log-concave. In particular,*

$$\eta_K^a(x)^c \eta_L^a(x)^{1-c} \le \eta_{cK+(1-c)L}^a(x) \le c\,\eta_K^a(x) + (1-c)\eta_L^a(x)$$

for $K, L, x \in (0,\infty)$ *and* $c \in (0,1)$, *with equality if and only if* $K = L$.

Proof. We observe that $f(K) = (s/s')^2$, where $s = \varphi_K^a(r)$ and $r = \sqrt{x/(x+1)}$. We get by Lemma 2.4(8)

$$f'(K) = \frac{8s^2\,\mathcal{K}_a(s)^2}{\pi^2 s'^2 K^2}\mu_a(r) = \frac{4}{\pi\sin(\pi a)}\frac{\mathcal{K}_a(r)}{\mathcal{K}'_a(r)}\left(\frac{s\,\mathcal{K}'_a(s)}{s'}\right)^2,$$

which is positive and increasing by Lemma 2.3(3), hence f is increasing and convex. For log-concavity, let $g(K) = \log(\eta_K^a(x))$. By Lemma 2.4(8) we get

$$g'(K) = \frac{8\,\mathcal{K}_a(s)^2}{\pi^2 K^2}\mu_a(r) = \frac{4}{\pi\sin(\pi a)}\frac{\mathcal{K}_a(r)}{\mathcal{K}'_a(r)}\mathcal{K}'_a(s)^2,$$

which is decreasing, hence f is log-concave. □

3.9. **Theorem.** *The function*

$$f(K) = \frac{\log\eta_K^a(x) - \log(x)}{K-1}$$

is decreasing from $(1,\infty)$ *onto*

$$\left(\frac{\pi\,\mathcal{K}_a(r)}{\sin(\pi\, a)\,\mathcal{K}'_a(r)}, \frac{4\,\mathcal{K}_a(r)\,\mathcal{K}'_a(r)}{\pi\sin(\pi\, a)}\right),$$

and the function

$$g(K) = \frac{\eta_K^a(x) - (x)}{K-1}$$

is increasing from $(1,\infty)$ *onto*

$$(4r^2\sin(\pi\, a)\,\mathcal{K}_a(r)\,\mathcal{K}'_a(r)/(\pi r'^2), \infty),$$

where $r = \sqrt{x/(x+1)}$.

Proof. It follows from Theorem 3.8 and Lemma 2.1 that f is monotone. Let $s = \varphi_K^a(r)$, by Lemma 2.4(6), the l'Hôpital Rule and definition of μ_a we get

$$\lim_{K\to1} f(K) = \lim_{K\to1}\frac{2}{K-1}\log\left(\frac{sr'}{s'r}\right)$$

$$= \lim_{K\to1}\frac{8\,\mathcal{K}_a(s)^2\mu_a(r)}{K^2\pi^2} = \frac{8}{\pi^2}\mathcal{K}_a(r)^2\mu_a(r) = \frac{4\,\mathcal{K}_a(r)\,\mathcal{K}'_a(r)}{\pi\sin(\pi\, a)}.$$

By using the fact that $K = \mu_a(r)/\mu_a(s)$ and the l'Hôpital Rule, we get

$$\lim_{K\to\infty} f(K) = \lim_{K\to\infty}\frac{8\mu_a(s)^2\,\mathcal{K}_a(s)^2}{\pi^2\mu_a(r)}$$

$$= \lim_{K\to\infty} \frac{2\,\mathcal{K}_a'(s)^2}{\sin^2(\pi\,a)\mu_a(r)} = \frac{2\,\mathcal{K}_a(0)^2}{\sin^2(\pi\,a)\mu_a(r)} = \frac{\pi\,\mathcal{K}_a(r)}{\sin(\pi\,a)\,\mathcal{K}_a'(r)}.$$

Next, let $g(K) = G(K)/H(K)$, where $G(K) = (s/s')^2 - (r/r')^2$ and $H(K) = K-1$. We see that $G(1) = H(1) = 0$ and $G(\infty) = H(\infty) = \infty$. We see that

$$G'(K)/H'(K) = 2(s\,\mathcal{K}_a'(s))^2/(s'^2\mu_a(r))\,,$$

and it follows from Lemma 2.3(3) and Lemma 2.1 that $g(K)$ is increasing and the required limiting values follow from $\varphi_K^a(r) = \mu_a^{-1}(\mu_a(r)/K)$. ☐

3.10. **Remark.** If we take $x = 1$ in Theorem 3.9, then with $t = 4\,\mathcal{K}_a(1/\sqrt{2})^2/(\pi\sin(\pi\,a))$ we have

(1) the function $\log(\lambda_a(K))/(K-1)$ is strictly decreasing from $(1,\infty)$ onto $(\pi/\sin(\pi\,a), t)$, and
(2) the function $(\lambda_a(K)-1)/(K-1)$ is increasing from $(1,\infty)$ onto $(t\,\sin^2(\pi\,a), \infty)$.

In particular,

$$e^{\pi(K-1)/\sin(\pi\,a)} < \lambda_a(K) < e^{t(K-1)},$$

$$1 + t(K-1)\sin^2(\pi\,a) < \lambda_a(K) < \infty,$$

respectively, and we get

$$\max\{e^{\pi(K-1)/\sin(\pi\,a)}, 1 + t(K-1)\sin^2(\pi\,a)\} < \lambda_a(K) < e^{t(K-1)}.$$

3.11. **Lemma.** *For $c \in [-3, 0)$, the function $f(r) = \mathcal{K}_a(r)^c + \mathcal{K}_a'(r)^c$ is strictly increasing from $(0, 1/\sqrt{2})$ onto $((\pi/2)^c, 2\,\mathcal{K}_a(1/\sqrt{2})^c)$.*

Proof. By Lemma 2.4(2) we get

$$f'(r) = \frac{2(1-a)c\,\mathcal{K}_a(r)^{c-1}\left(\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r)\right)}{rr'} - \frac{2(1-a)c\,\mathcal{K}_a'(r)^{c-1}(\,\mathcal{E}_a'(r) - r^2\,\mathcal{K}_a'(r))}{rr'}$$

$$= \frac{2(1-a)c(\,\mathcal{K}_a(r)\,\mathcal{K}_a'(r))^{c-1}}{rr'}(h(r) - h(r')),$$

and here $h(r) = \dfrac{r^2\,\mathcal{K}_a'(r)^{1-c}}{r^2}(\,\mathcal{E}_a(r) - r'^2\,\mathcal{K}_a(r))$, which is increasing on $(0,1)$ by [AVV1, Theorem 3.21(1)] and Lemma 2.3(3). Hence $f'(r) < 0$ on $(0, 1/\sqrt{2})$, and the limiting values are clear. ☐

3.12. **Theorem.** (1) *For $K > 1$, the function* $(\log(\lambda_a(K))/(K - 1/K)$ *is strictly increasing from* $(1,\infty)$ *onto* $(2\,\mathcal{K}_a(1/\sqrt{2})/(\pi\,\sin(\pi\,a)), \pi/\sin(\pi\,a))$.
(2) *The function* $\log(\lambda_a(K)+1)$ *is convex on* $(0,\infty)$*, and* $\log(\lambda_a(K))$ *is concave.*
(3) *The function* $g(K) = (\log(\lambda_a(K)))/\log K$ *is strictly increasing on* $(1,\infty)$*. In particular, for* $c \in (0,1)$

$$\lambda_a(K^c) < (\lambda_a(K))^c\,.$$

Proof. For (1), let $r = \mu_a^{-1}(\pi K/(2\sin(\pi\, a)))$, $0 \le r \le 1/\sqrt{2}$. Then by (1.3)

$$r' = \sqrt{1 - \left(\mu_a^{-1}\left(\frac{\pi K}{2\sin(\pi\, a)}\right)\right)^2}$$

$$= \sqrt{1 - \left(\mu_a^{-1}\left(K\mu_a\left(\frac{1}{\sqrt{2}}\right)\right)\right)^2} = \mu_a^{-1}\left(\frac{\pi}{2K\sin(\pi\, a)}\right),$$

we also observe that $K = \mathcal{K}'_a(r)/\mathcal{K}_a(r)$. Now it is enough to prove that the function

$$f(r) = \frac{2\log(r'/r)}{\mathcal{K}'_a(r)/\mathcal{K}_a(r) - \mathcal{K}_a(r)\mathcal{K}'_a(r)} = \frac{\pi\log(r'/r)}{\sin(\pi\, a)(\mu_a(r) + \mu_a(r'))},$$

is strictly decreasing on $(0, 1/\sqrt{2})$. Set $f(r) = G(r)/H(r)$. Clearly, $G(1/\sqrt{2}) = H(1/\sqrt{2}) = 0$. By Lemma 2.4(4) we get

$$\frac{G'(K)}{H'(K)} = \frac{4}{\pi\sin(\pi\, a)(\mathcal{K}_a(r)^{-2} - \mathcal{K}_a(r')^{-2})},$$

which is strictly decreasing from $(0, 1/\sqrt{2})$ onto

$$(2\mathcal{K}_a(1/\sqrt{2})/(\pi\ \sin(\pi\, a)), \pi/\sin(\pi\, a))$$

by Lemma 3.11. Now the proof of (1) follows from Lemma 2.1. For (2), it follows from Theorem 3.8 that $\log(\lambda_a(K))$ is concave. Letting $f(K) = \lambda_a(K) + 1$ we have

$$f(K) = \left(\mu_a^{-1}\left(\frac{\pi K}{2\sin(\pi\, a)}\right)\right)^{-2},$$

by (1.4) and (1.3). Now we have $\log f(K) = -2\log y$, here $\mu_a(y) = \pi K/(2\sin(\pi\, a))$. By Lemma 2.4(4) we get

$$\frac{f'(K)}{f(K)} = -\frac{2}{y}\frac{dy}{dK} = \frac{4}{\pi}(y'\,\mathcal{K}_a(y)),$$

which is decreasing in y by Lemma 2.3(3), and increasing in K. Hence $\log f(K)$ is convex.

For (3), $K > 1$, let $h(K) = (K - 1/K)/\log K$. We get

$$h'(K) = \frac{(1 + K^2)\log K - (K^2 - 1)}{(K\log K)^2},$$

which is positive because

$$\log K > \frac{2(K-1)}{K+1} > \frac{K^2 - 1}{K^2 + 1}$$

by [AVV1, 1.58(4)a], hence h is strictly increasing. Also

$$g(K) = h(K)\frac{\log(\lambda_a(K))}{K - 1/K} = \frac{\log(\lambda_a(K))}{\log K}$$

is strictly increasing by (1). This implies that

$$\frac{\log(\lambda_a(K^c))}{c\log K} < \frac{\log(\lambda_a(K))}{\log K},$$

and hence (3) follows. $\square$

3.13. **Corollary.** *For* $0<r<1/\sqrt{2}$ *and* $t=\pi^2/(2\,\mathcal{K}_a(1/\sqrt{2})^2)$*, we have*
(1) *The function* $f(r)=(\mu_a(r)-\mu_a(r'))/\log(r'/r)$ *is increasing from* $(0,1/\sqrt{2})$ *onto* $(1,t)$*. In particular,*

$$\log(r'/r) < \mu_a(r)-\mu_a(r') < \frac{\pi^2}{2\,\mathcal{K}_a(1/\sqrt{2})^2}\log(r'/r).$$

(2) *For* $g(r)=\log(r'/r)$*,*

$$g(r)+\sqrt{(\pi/\sin(\pi\, a))^2+g(r)^2} < 2\mu_a(r) < t\,g(r)+\sqrt{(\pi/\sin(\pi\, a))^2+t^2\,g(r)^2}.$$

Proof. It follows from the proof of Theorem 3.12(1) that $f(r)$ is increasing, and limiting values follows easily by the l'Hôpital Rule. For (2), from the definition of μ_a we get $\mu_a(r')=\pi^2/((2\sin(\pi\, a))^2\mu_a(r))$, replacing this in (1) we obtain

$$1<\frac{\mu_a(r)^2-\pi^2/(2\sin(\pi\, a))^2}{\mu_a(r)\log(r'/r)} < t = \frac{\pi^2}{2\,\mathcal{K}_a(1\sqrt{2})^2}\,.$$

This implies that

$$\mu_a(r)^2-\mu_a(r)\log(r'/r) > \frac{\pi^2}{(2\sin(\pi\, a))^2} \tag{3.14}$$

and

$$\mu_a(r)^2-t\,\mu_a(r)\log(r'/r) < \frac{\pi^2}{(2\sin(\pi\, a))^2}\,. \tag{3.15}$$

We get the left and right inequalities by solving (3.14) and (3.15) for $\mu_a(r)$, respectively. $\square$

4. THREE-PARAMETER COMPLETE ELLIPTIC INTEGRALS

The results in this section have counterpart in [AQVV]. For $a,b,c>0$, $a+b\geq c$, the decreasing homeomorphism $\mu_{a,b,c}:(0,1)\to(0,1)$, defined by

$$\mu_{a,b,c}(r)=\frac{B(a,b)}{2}\frac{F(a,b;c;r'^2)}{F(a,b;c;r^2)},\ r\in(0,1)$$

where B is the beta function. The (a,b,c)-modular function is defined by

$$\varphi_K^{a,b,c}(r)=\mu_{a,b,c}^{-1}(\mu_{a,b,c}(r)/K)\,.$$

We denote, in case $a<c$

$$\mu_{a,c}(r)=\mu_{a,c-a,c}(r) \quad \text{and} \quad \varphi_K^{a,c}(r)=\varphi_K^{a,c-a,c}(r)\,.$$

We define the three-parameter complete elliptic integrals of the first and second kinds for $0 < a < \min\{c, 1\}$ and $0 < b < c \le a + b$, by

$$\mathcal{K}_{a,b,c}(r) = \frac{B(a,b)}{2} F(a, b; c; r^2)$$

$$\mathcal{E}_{a,b,c}(r) = \frac{B(a,b)}{2} F(a-1, b; c; r^2)\,,$$

and denote

$$\mathcal{K}_{a,c}(r) = \mathcal{K}_{a,c-a,c}(r) \quad \text{and} \quad \mathcal{E}_{a,c}(r) = \mathcal{E}_{a,c-a,c}(r)\,.$$

4.1. **Lemma.** [HLVV, Theorem 3.6] *For* $0 < a < c \le 1$, *the function* $f(r) = \mu_{a,c}(r)\operatorname{artanh} r$ *is strictly increasing from* $(0, 1)$ *onto* $(0, (B/2)^2)$.

4.2. **Lemma.** [HLVV, Lemma 4.1] *Let* $a < c \le 1$, $K \in (1, \infty)$, $r \in (0, 1)$, *and let* $s = \varphi_K^{a,c}(r)$ *and* $t = \varphi_{1/K}^{a,c}(r)$. *Then the function*

(1) $f_1(r) = \mathcal{K}_{a,c}(s)/\mathcal{K}_{a,c}(r)$ *is increasing from* $(0, 1)$ *onto* $(1, K)$,
(2) $f_2(r) = s'\,\mathcal{K}_{a,c}(s)^2/(r'\,\mathcal{K}_{a,c}(r)^2)$ *is decreasing from* $(0, 1)$ *onto* $(0, 1)$,
(3) $f_3(r) = s\,\mathcal{K}'_{a,c}(s)^2/(r\,\mathcal{K}'_{a,c}(r)^2)$ *is decreasing from* $(0, 1)$ *onto* $(1, \infty)$,
(4) $g_1(r) = \mathcal{K}_{a,c}(t)/\mathcal{K}_{a,c}(r)$ *is decreasing from* $(0, 1)$ *onto* $(1/K, 1)$,
(5) $g_2(r) = t'\,\mathcal{K}_{a,c}(t)^2/(r'\,\mathcal{K}_{a,c}(r)^2)$ *is increasing from* $(0, 1)$ *onto* $(1, \infty)$,
(6) $g_3(r) = t\,\mathcal{K}'_{a,c}(t)^2/(r\,\mathcal{K}'_{a,c}(r)^2)$ *is increasing from* $(0, 1)$ *onto* $(0, 1)$,
(7) $g_4(r) = s/r$ *is decreasing from* $(0, 1)$ *onto* $(1, \infty)$,
(8) $g_5(r) = t/r$ *is increasing from* $(0, 1)$ *onto* $(0, 1)$.

4.3. **Theorem.** *For* $0 < a < c \le 1$, *the function* $f(x) = \mu_{a,c}(1/\cosh(x))$ *is increasing and concave from* $(0, \infty)$ *onto* $(0, \infty)$. *In particular,*

$$\mu_{a,c}\left(\frac{rs}{1 + r's'}\right) \le \mu_{a,c}(r) + \mu_{a,c}(s) \le 2\mu_{a,c}\left(\sqrt{\frac{2rs}{1 + rs + r's'}}\right)\,,$$

for all $r, s \in (0, 1)$. *The second inequality becomes equality if and only if* $r = s$.

Proof. Let $r = 1/\cosh(x)$ and (cf. [HLVV])

$$M(r^2) = \left(\frac{2}{B(a,b)}\right)^2 b\left(\mathcal{K}_{a,c}(r)\,\mathcal{E}'_{a,c}(r) + \mathcal{K}'_{a,c}(r)\,\mathcal{E}_{a,c}(r) - \mathcal{K}_{a,c}(r)\,\mathcal{K}'_{a,c}(r)\right).$$

We get

$$f'(x) = \frac{B(a,b)}{2}\frac{M(r^2)}{r'^2\,\mathcal{K}(r)^2}\,,$$

which is positive and increasing in r by [HLVV, Lemma 3.4(1), Theorem 3.12(2)], and f is decreasing in x. Hence f is concave. This implies that

$$\frac{1}{2}\left(\mu_{a,c}\left(\frac{1}{\cosh(x)}\right) + \mu_{a,c}\left(\frac{1}{\cosh(y)}\right)\right) \le \mu_{a,c}\left(\frac{1}{\cosh((x+y)/2)}\right)\,,$$

and we get the second inequality by using the formula

$$\left(\cosh\left(\frac{x+y}{2}\right)\right)^2=\frac{1+rs+r^{'}s^{'}}{2rs}$$

and setting $s=1/\cosh(y)$. Next, $f^{'}(x)$ is decreasing in x, and $f(0)=0$. Then $f(x)/x$ is decreasing on $(0,\infty)$ and $f(x+y)\le f(x)+f(y)$ by Lemmas 2.1 and 2.2, respectively. Hence the first inequality follows. $\square$

4.4. **Lemma.** *For* $0<a<c\le 1$*, we have*

$$\mu_{a,c}(r)+\mu_{a,c}(s)\le 2\mu_{a,c}(\sqrt{rs})\,,$$

for all $r,s\in(0,1)$*, with equality if and only if* $r=s$*.*

Proof. Clearly,

$$(r-s)^2\ge 0\Longleftrightarrow 1+r^2s^2\ge 1-(r-s)^2+r^2s^2$$

$$\Longleftrightarrow (1-rs)^2\ge 1-r^2-s^2+r^2s^2\Longleftrightarrow 1-rs\ge r^{'}s^{'}$$

$$\Longleftrightarrow 2\ge 1+rs+r^{'}s^{'}\Longleftrightarrow 1/(rs)\ge(1+rs+r^{'}s^{'})/(2rs)\,.$$

By using the fact that $\mu_{a,c}$ is decreasing, we get

$$\mu_{a,c}\left(\sqrt{\frac{2rs}{1+rs+r^{'}s^{'}}}\right)\le\mu_{a,c}(\sqrt{r\,s})\,,$$

and the result follows from Theorem 4.3. $\square$

4.5. **Theorem.** *For* $K>1$*,* $0<a<c$ *and* $r,s\in(0,1)$*,*

$$\tanh(K\operatorname{artanh}r)<\varphi_K^{a,c}(r).$$

The inequality is reversed if we replace K *by* $1/K$*.*

Proof. Let $s=\varphi_K^{a,c}(r)$. Then $s>r$, and by equality $\varphi_K^{a,c}(r)=\mu_{a,c}^{-1}(\mu_{a,c}(r)/K)$ and Lemma 4.1 we get

$$\frac{1}{K}\mu_{a,c}(r)\operatorname{artanh}s=\mu_{a,c}(s)\operatorname{artanh}s>\mu_{a,c}(r)\operatorname{artanh}r,$$

which is equivalent to the required inequality. For the case $1/K$ let $x=\varphi_{1/K}^{a,c}(r)$. Then $x<r$, and similarly we get

$$K\mu_{a,c}(r)\operatorname{artanh}x=\mu_{a,c}(x)\operatorname{artanh}x<\mu_{a,c}(r)\operatorname{artanh}r\,,$$

and this is equivalent to $\tanh((\operatorname{artanh}r)/K)>\varphi_{1/K}^{a,c}(r)$. $\square$

References

[AS] M. Abramowitz, I. Stegun: *Handbook of mathematical functions with formulas, graphs and mathematical tables.* National Bureau of Standards, 1964 (Russian translation, Nauka 1979).

[AQ] H. Alzer and S.-L. Qiu: *Monotonicity theorems and inequalities for the complete elliptic integrals.* J. Comput. Appl. Math. 172 (2004), no. 2, 289–312.

[AQVV] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy, and M. Vuorinen: *Generalized elliptic integrals and modular equation.* Pacific J. Math. Vol. 192, No. 1, (2000), 1–37.

[AVV1] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Conformal invariants, inequalities and quasiconformal maps.* J. Wiley, 1997, 505 pp.

[AVV2] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Dimension-free quasiconformal distortion in n-space.* Trans. Amer. Math. Soc. 297 (1986), 687–706.

[AVV3] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Inequalities for the extremal distortion function.* - Proc. of the 13th Rolf Nevanlinna Colloquium, Lecture Notes in Math. Vol. 1351, 1–11, Springer-Verlag, 1988

[AVV4] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Topics in special functions.* - Papers on Analysis: A volume dedicated to Olli Martio on the occasion of his 60th birthday, ed. by J. Heinonen, T. Kilpeläinen, and P. Koskela, Report 83, Univ. Jyväskylä (2001), 5–26, ISBN 951-39-1120-9, arXiv:0712.3856.

[AVV5] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Topics in special functions II.* - Conform. Geom. Dyn. 11 (2007), 250–270.

[AV] G. D. Anderson and M. Vuorinen: *Reflections on Ramanujan's Mathematical Gems.* Mathematics Newsletter: Special Issue Commemorating ICM 2010 in India Vol. 19, Sp. No. 1, August 2010, pp. 87108, http://www.ramanujanmathsociety.org/mnl/August2010/mnl-spl-fullfiles.pdf

[BPV] R. Balasubramanian, S. Ponnusamy and M. Vuorinen: *Functional inequalities for quotients of hepergeometric functions.* J. Math. Anal. Appl. 218 (1998) 256–268.

[B1] Á. Baricz: *Functional inequalities involving special functions.* J. Math. Anal. Appl. 319(2)(2006) 450–459.

[B2] Á. Baricz: *Functional inequalities involving special functions II.* J. Math. Anal. Appl. 327(2)(2007) 1202–1213.

[B3] Á. Baricz: *Turán type inequalities for generalized complete elliptic integrals.* Math. Z. 256(4)(2007) 895–911.

[B4] Á. Baricz: *Turán type inequalities for hypergeometric functions.* Proc. Amer. Math. Soc. 136 (2008), no. 9, 3223–3229.

[B5] Á. Baricz: *Generalized Bessel functions of the first kind.* Lecture Notes in Mathematics, 1994. Springer-Verlag, Berlin, 2010. xiv+206 pp.

[BBG] B. C. Berndt, S. Bhargava, and F. G. Garvan: *Ramanujan's theories of elliptic functions to alternative bases.* Trans. Amer. Math. Soc. 347 (1995), no. 11, 4163–4244.

[BV] B. A. Bhayo and M. Vuorinen: *Inequalities for eigenfunctions of the p-Laplacian.* arXiv:1101.3911[math.CA].

[HRV] H. Hakula, A. Rasila, and M. Vuorinen: *On moduli of rings and quadrilaterals: algorithms and experiments.*- SIAM J. Sci. Comput. 33 (2011) 279–302, arXiv:0906.1261 [math.NA].

[HLVV] V. Heikkala, H. Lindén, M. K. Vamanamurthy, and M. Vuorinen: *Generalized elliptic integrals and the Legendre M-function.* J. Math. Anal. Appl. 338 (2008) 223–243, arXiv:math/0701438.

[HVV] V. Heikkala, M. K. Vamanamurthy, and M. Vuorinen: *Generalized elliptic integrals.* Comput. Methods Funct. Theory 9 (2009), no. 1, 75–109, arXiv:math/0701436.

[K] R. Kühnau, ed.: Handbook of complex analysis: geometric function theory. Vol. 1-2. North-Holland, Amsterdam, 2002 and 2005.

[LV] O. Lehto and K. I. Virtanen: *Quasiconformal mappings in the plane.* Second edition. Translated from the German by K. W. Lucas. Die Grundlehren der mathematischen Wissenschaften, Band 126. Springer-Verlag, New York-Heidelberg, 1973. viii+258 pp.

[L] P. Lindqvist: *Some remarkable sine and cosine functions.* Ricerche di Matematica, Vol. XLIV (1995), 269-290.

[PV] S. Ponnusamy and M. Vuorinen: *Asymptotic expansions inequalities for hypergeometric functions.* Mathematika 44 (1997), 278–301.

[QV] S.-L. Qiu and M. Vuorinen: *Duplication inequalities for the ratios of hypergeometric functions.* Forum Math. 12(1)(2000), 109–133.

[Ru] H. Ruskeepää: *Mathematica® Navigator.* 3rd ed. Academic Press, 2009.

[Vu1] M. Vuorinen: *Conformal geometry and quasiregular mappings*, Lecture Notes in Mathematics 1319, Springer, Berlin, 1988.

[Vu2] M. Vuorinen: *Special functions and conformal invariants.* Analysis and its applications (Chennai, 2000), 255–268, Allied Publ., New Delhi, 2001.

[WZC] G. Wang, X. Zhang, and Y. Chu: *Inequalities for the generalized elliptic integrals and modular functions.* J. Math. Anal. Appl. 331 (2007), no. 2, 1275–1283.

[WZQC] G. Wang, X. Zhang, S. Qiu, and Y. Chu: *The bounds of the solutions to generalized modular equations.* J. Math. Anal. Appl. 321 (2006), no. 2, 589–594.

[ZWC1] X. Zhang, G. Wang, and Y. Chu: *Some inequalities for the generalized Grötzsch function.* Proc. Edinb. Math. Soc. (2) 51 (2008), no. 1, 265–272.

[ZWC2] X. Zhang, G. Wang, and Y. Chu: *Remarks on generalized elliptic integrals.* Proc. Roy. Soc. Edinburgh Sect. A 139 (2009), no. 2, 417–426.

Department of Mathematics, University of Turku, FI-20014 Turku, Finland
E-mail address: `barbha@utu.fi`

Department of Mathematics, University of Turku, FI-20014 Turku, Finland
E-mail address: `vuorinen@utu.fi`